arXiv:0903.5090v1 [math.DG] 29 Mar 2009

MINIMAL SURFACES IN QUASI-FUCHSIAN 3-MANIFOLDS

Biao Wang

Abstract

In this paper, we prove that if a quasi-Fuchsian 3-manifold M contains a closed geodesic with complex length $\mathscr{L} = l + i\theta$ such that $\theta/l \gg 1$, then it contains at least two minimal surfaces which are incompressible in M.

1. Introduction

For any complete hyperbolic 3-manifold N, there is a holonomy representation $\rho : \pi_1(N) \to \mathrm{Isom}^+(\mathbb{H}^3)$ so that $G := \rho(\pi_1(N))$ is a discrete subgroup of $\mathrm{Isom}^+(\mathbb{H}^3)$ called *Kleinian group* and $N = \mathbb{H}^3/G$. For any torsion free Kleinian group $G \subset \mathrm{Isom}^+(\mathbb{H}^3)$, the quotient $\mathbb{H}^3/G$ is a complete hyperbolic 3-manifold.

For a torsion free Kleinian group G, if its limits set $\Lambda(G) \subset S^2_\infty$ is a closed Jordan curve, i.e. $\Lambda(G)$ is homeomorphic to the unit circle S^1, then G is called a *quasi-Fuchsian group*, and $M = \mathbb{H}^3/G$ is called a *quasi-Fuchsian* 3*-manifold*. The quasi-Fuchsian 3-manifold M has a *convex core*, denoted by $\mathscr{C}(M)$, which is the quotient of the convex hull of $\Lambda(G)$. Topologically, M is homeomorphic to $\Sigma \times \mathbb{R}$, where Σ is a finite type Riemann surface with negative Euler characteristic. In this paper, Σ is assumed to a closed surface with genus $\geqslant 2$ except in section 4 where Σ denotes a hypersurface in a Riemannian manifold.

Any quasi-Fuchsian 3-manifold $M = \Sigma \times \mathbb{R}$ always contains a minimal surface, say T, which is incompressible in M (cf. [**And83**, **SY79**, **Uhl83**]). Besides, K. Uhlenbeck also proved that if the principle curvature of T is between -1 and 1, then $T \subset M$ is unique and M can be foliated by equidistant surfaces to T (cf. [**Uhl83**]). In this case, we proved that M admits a foliation such that each leaf is a surface of constant mean curvature (cf. [**Wan08**]).

In this paper, we will prove the following theorem.

Mathematics Subject Classification. Primary 53A10, Secondary 57M05.

Received March 28, 2009.

Theorem 1.1. *Let* $M = \Sigma \times \mathbb{R}$ *be a quasi-Fuchsian* 3*-manifold, let* $\gamma \subset M$ *be a simple closed geodesics* γ *with complex length* $\mathscr{L} = l + i\theta$*, and let* $S \subset M \setminus \gamma$ *be an embedded closed surface that is incompressible in* M*. If* $\theta/l \gg 1$*, then there exists a least area minimal surface* $T \subset M \setminus \gamma$ *that is isotopic to* S *in* $M \setminus \gamma$*, here* T *is of least area means if there is another minimal surface* $T' \subset M \setminus \gamma$ *isotopic to* S*, then* $\operatorname{Area}(T) \leqslant \operatorname{Area}(T')$*.*

If a quasi-Fuchsian 3-manifold contains more than one sufficiently short simple closed geodesics, say $\gamma_1, \dots, \gamma_n$, then we have a result similar to Theorem 1.1, see Corollary 6.2 for detail.

The plan of this paper is as follows. In section 2 we review the basic properties of tubes, in section 3 we discuss some properties of helicoids in $\mathbb{H}^3$, in section 4 we prove one version of coarea formulae used frequently in [**CG06**], and in section 5 we introduce a tool called shrinkwrapping developed by Calegari and Gabai in [**CG06**], in section 6 we prove Theorem 1.1.

Acknowledgement. I would like to express my gratitude to my advisor Profesoor Bill Thurston for his help, stimulating suggestions and encouragement in the past several years.

2. Tubes of short geodesics

We consider the upper half space model of hyperbolic 3-space, that is, a three-diemnsional space

$$\mathbb{H}^3 = \{z + tj \mid z \in \mathbb{C},\ t \in \mathbb{R},\ t > 0\}$$

which is equipped with the (hyperbolic) metric

$$ds^2 = \frac{|dz|^2 + dt^2}{t^2}\ ,$$

where $z = x + iy$ for $x, y \in \mathbb{R}$. Let $\mathrm{PSL}_2(\mathbb{C})$ denote the set of Möbius transformations on $\mathbb{H}^3$. For any $g \in \mathrm{PSL}_2(\mathbb{C})$, write

$$g(z) = \frac{az + b}{cz + d}\ , \quad \forall\, z \in \mathbb{C}\ ,$$

then the Poincaré extension of g is given by

$$g(z + tj) = \frac{(az + b)(\overline{cz + d}) + a\bar{c}t^2 + tj}{|cz + d|^2 + |c|^2 t^2}\ . \tag{2.1}$$

In this paper, we are interested in the loxodromic elements, i.e. by conjugacy, $g(z) = \alpha z$, where $\alpha = \exp(l + i\theta)$ with $l > 0$ and $0 \leqslant \theta < 2\pi$. The loxodromic transformation

$$g(z + tj) = \alpha z + |\alpha| tj \tag{2.2}$$

translates points on the t-axis by distance l and twists a normal plane by angle θ. The complex number $\mathscr{L} := l + i\theta$ is called the *complex length* of g. Since the trace of g is invariant under conjugate transformations, the complex length of g is also invariant under conjugate transformations. Notice that g determines θ only up to a multiple of 2π.

Now suppose γ is a simple closed geodesic in a complete hyperbolic 3-manifold N, then a lift $\tilde{\gamma}$ in $\mathbb{H}^3$ is the axis of a loxodromic element $g \in \mathrm{PSL}_2(\mathbb{C})$ representing γ. The choice of different lift $\tilde{\gamma}'$ gives rise to another element of $\mathrm{PSL}_2(\mathbb{C})$ such that two elements are conjugate in $\mathrm{PSL}_2(\mathbb{C})$. Two lifts $\tilde{\gamma}$ and $\tilde{\gamma}'$ possesses a common perpendicular. Let $r_{\max}(\gamma)$ be the maximal number $r > 0$ such that $\mathscr{N}_r(\tilde{\gamma}) \cap \mathscr{N}_r(\tilde{\gamma}') = \emptyset$ as $\tilde{\gamma}'$ runs through all the lifts of γ different from $\tilde{\gamma}$, where $\mathscr{N}_r(\tilde{\gamma}) = \{x \in \mathbb{H}^3 \mid \mathrm{dist}(x, \tilde{\gamma}) < r\}$ is the r-neighborhood of the geodesic $\tilde{\gamma}$ in $\mathbb{H}^3$. We call $r_{\max} = r_{\max}(\gamma)$ the *tube radius* of γ. The (maximal solid) tube of γ is defined by

$$\mathbb{T}(\gamma) = \mathscr{N}_{r_{\max}}(\tilde{\gamma})/\langle g \rangle \ . \tag{2.3}$$

When l, the real length of γ, is sufficiently small, R. Meyerhoff found a relation between $r_{\max}$ and l.

Theorem 2.1 ([**Mey87**]). *If γ is a closed geodesic in a complete hyperbolic 3-manifold N, with complex length $\mathscr{L} = l + i\theta$ satisfying*

$$l \leqslant \frac{\sqrt{3}}{4\pi}\,[\log(\sqrt{2}+1)]^2 \approx 0.107 \ , \tag{2.4}$$

then there exists an embedded solid tube around γ whose radius is given by the following formula

$$\sinh^2 r_{\max}(\gamma) = \frac{1}{2}\left(\frac{\sqrt{1-2\kappa(l)}}{\kappa(l)} - 1\right) \ , \tag{2.5}$$

where $\kappa(l) = \cosh(\sqrt{4\pi l}/\sqrt[4]{3}) - 1$.

If a hyperbolic 3-manifold N contains more than one short geodesics, we want the solid tubes of different geodesics to be disjoint. Fortunately, R. Meyerhoff also proved the following theorem.

Theorem 2.2 ([**Mey87**]). *The solid tubes constructed by* (2.3) *about different short geodesics whose real length satisfy* (2.4) *do not intersect.*

3. Helicoids in $\mathbb{H}^3$

In this section, we consider the Lorentzian 4-space $\mathbb{L}^4$, i.e. a vector space $\mathbb{R}^4$ with the Lorentzian inner product

$$\langle x\,, y\rangle = -x_1y_1 + x_2y_2 + x_3y_3 + x_4y_4$$

where $x, y \in \mathbb{R}^4$. The hyperbolic space $\mathbb{H}^3$ can be considered as the unit sphere of $\mathbb{L}^4$:

$$\mathbb{H}^3 = \{x \in \mathbb{L}^4 \mid \langle x\,, x\rangle = -1,\, x_1 \geqslant 1\}\ .$$

The helicoid $\mathcal{H}_a$ is the surface parametrized by the (u, v)-plane in the following way:

$$\mathcal{H}_a = \left\{ x \in \mathbb{H}^3 \;\middle|\; \begin{array}{ll} x_1 = \cosh u \cosh v, & x_2 = \cosh u \sinh v \\ x_3 = \sinh u \cos(av), & x_4 = \sinh u \sin(av) \end{array} \right\} \tag{3.1}$$

The axis of the helicoid $\mathcal{H}_a$ is

$$(\cosh v, \sinh v, 0, 0)\ , \qquad -\infty < v < \infty\ ,$$

which is the intersection of x_1x_2-plane and the three dimensional hyperboloid. The first fundamental form of $\mathcal{H}_a$ is

$$I = du^2 + (\cosh^2 u + a^2 \sinh^2 u)dv^2\ . \tag{3.2}$$

Intuitively, when $a > 0$ is sufficiently large, the helicoid $\mathcal{H}_a$ will become unstable. Actually, Hiroshi Mori proved the following theorem.

Theorem 3.1 ([**Mor82**]). *Let $\mathcal{H}_a$ be the helicoid as above, then it is a minimal surface in $\mathbb{H}^3$. Furthermore,*

- *$\mathcal{H}_a$ is globally stable if $a \leqslant 3\sqrt{2}/4$,*
- *$\mathcal{H}_a$ is globally unstable if $a \geqslant \sqrt{105\pi}/8$.*

The unstability of $\mathcal{H}_a$ when a is sufficiently large is very important in this paper. We also need the following lemma, which shows that $\mathcal{H}_a$ is a ruled surface in $\mathbb{H}^3$.

Lemma 3.2 (Tuzhilin [**Tuz93**]). *Let $\mathcal{H}_a$ be the helicoid as above, then it is a ruled surface, i.e. it is stratified into straight lines in the sense of hyperbolic metric.*

Let γ be a closed geodesic in a hyperbolic 3-manifold N, let $g \in \mathrm{PSL}_2(\mathbb{C})$ be a loxodromic element with complex length $\mathscr{L} = l + i\theta$ which represents γ, and let $\tilde{\gamma} \subset \mathbb{H}^3$ be the axis of g. Without of lose generality, we may assume $\theta > 0$. Write $a = \theta/l$, let $\mathcal{H}_{\theta/l} \subset \mathbb{H}^3$ be the helicoid constructed as above, and let

$$\mathcal{A}_{\theta/l} = \frac{\mathcal{H}_{\theta/l} \cap \mathscr{N}_{r_{\max}}(\tilde{\gamma})}{\langle g\rangle}\ . \tag{3.3}$$

Lemma 3.3. *The quotient surface $\mathcal{A}_{\theta/l}$ is a minimal annulus in the tube $\mathbb{T}(\gamma)$, and its area is*

$$\operatorname{Area}(\mathcal{A}_{\theta/l}) = 2\int_0^{r_{\max}} (l^2 \cosh^2 u + \theta^2 \sinh^2 u)^{1/2} du\ . \tag{3.4}$$

Proof. For each point $p \in \mathcal{H}_{\theta/l}$, the point $g(p) \in \mathcal{H}_{\theta/l}$ is obtained by twisting by angle θ and translating by distance l, so $\mathcal{H}_{\theta/l}$ is invariant under the cyclic group $\langle g \rangle$, the quotient surface $\mathcal{A}_{\theta/l}$ is an annulus in $\mathbb{T}(\gamma)$. By Lemma 3.2 and coarea formula (4.2), the area of $\mathcal{A}_{\theta/l}$ can be computed as follows:

$$\begin{aligned}\mathrm{Area}(\mathcal{A}_{\theta/l}) &= \int_0^{r_{\max}} \mathrm{Length}(\mathcal{A}_{\theta/l} \cap \mathscr{N}_r(\gamma)) dr \\ &= 2\int_0^{r_{\max}} \left\{ \int_0^l \left[\cosh^2 r + \left(\frac{\theta}{l}\right)^2 \sinh^2 r \right]^{1/2} ds \right\} dr \\ &= 2\int_0^{r_{\max}} (l^2 \cosh^2 r + \theta^2 \sinh^2 r)^{1/2} dr \ .\end{aligned}$$

Therefore the lemma is proved. □

Theorem 3.4. *Suppose a closed geodesic γ in a complete hyperbolic 3-manifold N has complex length $\mathscr{L} = l + i\theta$, let $\mathbb{T}(\gamma)$ and $\mathcal{A}_{\theta/l}$ be constructed as above. If θ/l is sufficiently large, then*

$$\mathrm{Area}(\mathcal{A}_{\theta/l}) > \mathrm{Area}(\partial\mathbb{T}(\gamma)) \ .$$

Proof. By Lemma 3.3, when $\theta/l \gg 1$, we have the following estimate

$$\mathrm{Area}(\mathcal{A}_{\theta/l}) \geqslant 2\theta \int_0^{r_{\max}} \sinh u \, du = 2\theta(\cosh r_{\max} - 1) \ .$$

Since $r_{\max}(\gamma) \to \infty$ as $l \to 0$, we have the estimate

$$\mathrm{Area}(\mathcal{A}_{\theta/l}) \geqslant \theta \cosh r_{\max}$$

if $l \ll 1$. On the other hand, the area of the torus $\partial\mathbb{T}(\gamma)$ is given by

$$\mathrm{Area}(\partial\mathbb{T}(\gamma)) = 2\pi l \cosh r_{\max} \sinh r_{\max} \ .$$

As $l \to 0$, we claim that $l \sinh r_{\max} \to 0$. In fact, by Theorem 2.1, we have the identity

$$l^2 \sinh^2(r_{\max}) = \frac{l^2}{2}\left(\frac{\sqrt{1-2\kappa(l)}}{\kappa(l)} - 1\right) \ ,$$

where $\kappa(l) = \cosh(\sqrt{4\pi l}/\sqrt[4]{3}) - 1 = 2\pi l/\sqrt{3} + \mathcal{O}(l^2)$. Therefore

$$l^2 \sinh^2(r_{\max}) = \frac{l}{2\pi/\sqrt{3} + \mathcal{O}(l)} \cdot \frac{\sqrt{1-2\kappa(l)} - \kappa(l)}{2} \to 0 \ ,$$

as $l \to 0$. Therefore when $\theta/l \gg 1$, we have the estimate $2\pi l \sinh r_{\max} < \theta$, which implies the lemma. □

Remark 1. The area of the meridian disk of $\mathbb{T}(\gamma)$ is $2\pi(\cosh r_{\max}-1)$, so the computation in Theorem 3.4 also implies that the area of the meridian disk is bigger than $\text{Area}(\partial\mathbb{T}(\gamma))$ when θ/l is sufficiently large.

Remark 2. From the computation above, as $l \to 0$, we have

$$\text{Area}(\partial\mathbb{T}(\gamma)) \to \frac{\sqrt{3}}{2} \qquad \text{and} \qquad \text{Vol}(\mathbb{T}(\gamma)) \to \frac{\sqrt{3}}{4}\ .$$

4. Coarea formula

The following coarea formulae (4.1) and (4.2) were used frequently in [**CG06**] without proof. Since the coarea formulae are very important in this paper too, we give a proof here for convenience, which is also pretty simple.

Lemma 4.1. *Suppose Σ is a hypersurface in a complete Riemannian manifold $(N, \bar{g}_{ij})$. Let $p \in N$ be a fixed point, for any point $q \in \Sigma$, define $\alpha(q)$ to be the angle between the tangent space to Σ at q, and the radial geodesic through q emanating from p. Then*

$$\text{Area}(\Sigma \cap \mathscr{N}_s(p)) = \int_0^s \int_{\Sigma\cap\partial\mathscr{N}_t(p)} \frac{1}{\cos\alpha}\, dldt\ . \tag{4.1}$$

Proof. Define a function $h : \Sigma \to \mathbb{R}$ by $h(q) = \text{dist}(p, q)$, where $\text{dist}(\cdot,\cdot)$ is the distance function on N. Then, by the standard coarea formula (cf. [**SY94**, p. 89]), we have

$$\text{Area}(\Sigma \cap \mathscr{N}_s(p)) = \int_0^s \int_{\Sigma\cap\partial\mathscr{N}_t(p)} \frac{1}{|\nabla_\Sigma h|}\, dldt\ .$$

Since $\nabla h = (\nabla h)^\perp + (\nabla h)^\parallel = \nabla^\perp h + \nabla_\Sigma h$ and $|\nabla h|_{\bar{g}_{ij}} = 1$, one has $|\nabla_\Sigma h| = \cos\alpha$. □

Remark 3. If γ is a simple closed geodesic in N, then we still have the following coarea formula

$$\text{Area}(\Sigma \cap \mathscr{N}_s(\gamma)) = \int_0^s \int_{\Sigma\cap\partial\mathscr{N}_t(\gamma)} \frac{1}{\cos\alpha}\, dldt\ . \tag{4.2}$$

5. Shrinkwrapping

In this section, we use a technical tool, called shrinkwrapping developed by Calegari and Gabai in [**CG06**], to find minimal surfaces in quasi-Fuchsian 3-manifolds containing a very short simple closed geodesic. Roughly speaking, given a simple closed geodesic γ in a quasi-Fuchsian 3-manifold M and an embedded surface $S \subset M \setminus \gamma$, a surface $T \subset M$ is obtained from S by *shrinkwrapping S rel. γ* if it homotopic to S, can be approximated by an isotopy from S supported in $M \setminus \gamma$, and is least area subject to these constraints.

Definition 5.1. Choose some positive σ so that $\mathscr{N}_\sigma(\gamma) \subset M$ is an embedded solid torus , for $0 \leqslant t < 1$ we define a family of Riemannian metrics (M, g_t) in the following manner.

Let $h : \mathscr{N}_{\sigma(1-t)}(\gamma) \to [0, \sigma(1-t)]$ be the function whose value at a point p is the hyperbolic distance from p to γ. Let $\phi : [0,1] \to [0,1]$ be a C^∞ bump function given by

$$\phi(t) = \begin{cases} 0, & t \in [0, 1/4], \\ \text{strictly increasing}, & t \in [1/4, 1/3], \\ 1, & t \in [1/3, 2/3], \\ \text{strictly decreasing}, & t \in [2/3, 3/4], \\ 0, & t \in [3/4, 1]\ . \end{cases}$$

We define a metric g_t on M:

- g_t agrees with the hyperbolic metric on $M \setminus \mathscr{N}_{\sigma(1-t)}(\gamma)$;
- g_t is conformally equivalent to the hyperbolic metric on $\mathscr{N}_{\sigma(1-t)}(\gamma)$, which is more precisely given by the ratio

$$\frac{g_t \text{ length element}}{\text{hyperbolic length element}} = 1 + 2\phi\left(\frac{h(p)}{\sigma(1-t)}\right) .$$

Lemma 5.2 ([**CG06**])**.** *The g_t metric satisfies the following properties:*

(i) *For each t there is an $f(t)$ satisfying $\frac{2}{3}\sigma(1-t) < f(t) < \frac{3}{4}\sigma(1-t)$ such that the union of tori $\partial\mathscr{N}_{f(t)}(\gamma)$ are totally geodesic for the g_t metric.*

(ii) *The restricted metric $g_t|\mathscr{N}_\sigma(\gamma)$ admits a family of isometries preserving γ and acts transitively on the unit normal bundle (in M) to γ.*

(iii) *The area of a disk cross–section on $\mathscr{N}_{\sigma(1-t)}$ is $O((1-t)^2)$.*

(iv) *The metric g_t dominates the hyperbolic metric on 2-planes. That is, for all 2-vectors ν, the g_t area of ν is at least as large as the hyperbolic area of ν.*

Lemma 5.3 ([**CG06**])**.** *Let M, γ and S be as in the statement of Theorem* 1.1*. Let $f(t)$ be a function given in Lemma* 5.2 *so that $\partial\mathscr{N}_{f(t)}(\gamma)$ is totally geodesic with respect to the g_t metric. Then for each t, there exists an embedded surface S_t isotopic in $M \setminus \mathscr{N}_{f(t)}(\gamma)$ to S, and which is globally g_t-least area among all such surfaces.*

Proof. By Lemma 5.2, the surfaces $\partial\mathscr{N}_{f(t)}(\gamma)$ are totally geodesic with respect to the g_t metrics, therefore these surfaces can act as barrier surfaces. As a quasi-Fuchsian 3-manifold, M has a compact convex core $\mathscr{C}(M)$ whose boundary consists of two incompressible surfaces which are convex with respect to the inward normal. Let

$$M_t = \mathscr{C}(M) \setminus \mathscr{N}_{f(t)}(\gamma) \ ,$$

then M_t is a compact manifold with mean convex boundary. We assume, after a small isotopy if necessary, that S does not intersect $\mathscr{N}_{f(t)}(\gamma)$ for any $t \in [0,1]$, therefore S is an incompressible surface in M_t. Then Meeks-Simon-Yau's main theorem in [**MSY82**] implies that there exists a globally least area surface S_t with respect to the metric g_t. □

6. Proof of Theorem 1.1

Lemma 6.1. *Suppose γ is a closed geodesic in a complete hyperbolic 3-manifold M, let C be a simple null-homotoptic curve on $\partial\mathscr{N}_r(\gamma)$ with* $\mathrm{Length}(C) < 2\pi\sinh r$*, here $r > 0$ is chosen so that the r-tube $\mathscr{N}_r(\gamma)$ is embedded, then C bounds an embedded least area disk-type minimal surface $\Delta \subset \mathscr{N}_r(\gamma) \setminus \gamma$.*

Proof. Suppose C bounds a disk $D \subset \partial\mathscr{N}_r(\gamma)$, then the diameter of D, denoted by $\mathrm{diam}(D)$, with respect to the induced metric on $\partial\mathscr{N}_r(\gamma)$ is $< \pi\sinh r$; otherwise we have $\mathrm{Length}(C) \geqslant 2\pi\sinh r$, which implies a contradiction.

Since the longitudes $\mathcal{A}_{\theta/l} \cap \partial\mathscr{N}_r(\gamma)$ divide the torus $\partial\mathscr{N}_r(\gamma)$ into two semi-tori $R_\pm$, and recall that the length of the meridian of $\partial\mathscr{N}_r(\gamma)$ is $2\pi\sinh r$, so D must be contained in one of the semi-tori, say R_+. These longitudes bound a minimal annulus $\mathcal{A}_{\theta/l}(r) := \mathcal{A}_{\theta/l} \cap \mathscr{N}_r(\gamma)$, then $R_+ \cup \mathcal{A}_{\theta/l}(r)$ is mean convex with respect to the inward normal, by a result in [**MY82a**, **MY82b**], we know that C bounds a least area disk Δ contained in the semi-tube bounded by R_+ and $\mathcal{A}_{\theta/l}(r)$, and any such least area disk is properly embedded. The minimal disk Δ cannot intersect $\mathcal{A}_{\theta/l}(r)$, since the latter is a barrier surface. In particular, Δ must be disjoint from the closed geodesic $\gamma \subset \mathcal{A}_{\theta/l}(r)$. □

Remark 4. If γ is very short and $r \in [r_{\max}/4, r_{\max}/2]$, where $r_{\max}$ is the tube radius of γ, then Lemma 6.1 can be proved without using Meeks-Yau's result.

- The existence of Δ is from J. Douglas, T. Radó and C. B. Morrey.
- If Δ intersects γ, then Δ contains an intrinsic disk $\mathcal{B}_r$ of radius r, and $\mathrm{Area}(\mathcal{B}_r) \geqslant 2\pi(\cosh r - 1)$. Since D must be contained

in a semi-torus, we have $\mathrm{Area}(D) \leqslant \pi l \sinh r \cosh r$. Therefore $2\pi(\cosh r - 1) \leqslant \pi l \sinh r \cosh r$. This is impossible when $r_{\max}$ is sufficiently large.
- If $\Delta \not\subset \mathscr{N}_r(\gamma)$, then the same argument as above tells us that $\Delta \subset \mathscr{N}_{2r}(\gamma)$, and so there exists $r_0 \in (r, 2r)$ such that $\partial\mathscr{N}_{r_0}(\gamma)$ touches Δ for the first time from outside, but this is impossible by maximum principle.

Therefore C bounds a least area minimal disk $\Delta \subset \mathscr{N}_r(\gamma) \setminus \gamma$.

Now we can prove Theorem 1.1.

Proof of Theorem 1.1. Let $\sigma < r_{\max}/2$ be a constant, where $r_{\max}$ is the tube radius of γ that will be determined later. By Lemma 5.3, we can construct g_t-least area minimal surface S_t that is isotopic to S for $t \in [0,1)$.

Claim. *Suppose $\theta/l \gg 1$, then there exists a constant $\varepsilon = \varepsilon(l,\theta)$ such that $S_t \cap \mathscr{N}_\varepsilon(\gamma) = \emptyset$ for all t that is sufficiently large.*

Proof of Claim. At first, by Lemma 6.1, there exists a constant $\varepsilon > 0$ depending only on $r_{\max}$ such that for any simple loop $C \subset \partial\mathscr{N}_r(\gamma)$ with $\mathrm{Length}(C) < 2\pi \sinh r$, each least area disk $\Delta \subset \mathscr{N}_r(\gamma)$ bounded C is ε away from γ, i.e. $\mathrm{dist}(\Delta, \gamma) \geqslant \varepsilon$, where $r \in [r_{\max}/2, r_{\max}]$. Next choose t sufficiently large such that $\sigma(1-t) < \varepsilon$.

For any $t \in [0,1)$, $S_t \cap \mathbb{T}(\gamma)$ is either empty or the union of annuli and disks, otherwise S_t would be not incompressible. For any sufficiently large $t \in [0,1)$, if $S_t \cap \mathscr{N}_{r_{\max}/2}(\gamma) = \emptyset$, then we are done; so we assume

$$S_t \cap \mathscr{N}_{r_{\max}/2}(\gamma) \neq \emptyset \ , \quad \text{for all large } t \in [0,1) \ .$$

We don't need worry about the null-homotopic loops on $\partial\mathscr{N}_r(\gamma)$ with (hyperbolic) length $< 2\pi \sinh r$ for all $r \in [r_{\max}/2, r_{\max}]$. In fact, suppose $\Delta \subset S_t \cap \mathscr{N}_r(\gamma)$ is a disk such that $\mathrm{Length}(\partial\Delta) < 2\pi \sinh r$, then $\partial\Delta$ bounds a least area minimal surface $\Delta' \subset \mathscr{N}_r(\gamma) \setminus \mathscr{N}_\varepsilon(\gamma)$ with respect to the hyperbolic metric, which is also a g_t-least area disk since $\sigma(1-t) < \varepsilon$, this implies $\Delta = \Delta'$. So Δ is disjoint from $\mathscr{N}_\varepsilon(\gamma)$.

Now we can assume that for any $r \in [r_{\max}/2, r_{\max}]$, the boundary of each component of $S_t \cap \mathscr{N}_r(\gamma)$ which is a disk has length $> 2\pi \sinh r$. By Lemma 5.2 and coarea formula (4.2), we have estimate

$$\begin{aligned}
\mathrm{Area}(S_t \cap \mathbb{T}(\gamma), g_t) &\geqslant \mathrm{Area}(S_t \cap \mathbb{T}(\gamma)) \\
&\geqslant \mathrm{Area}(S_t \cap (\mathbb{T}(\gamma) \setminus \mathscr{N}_{r_{\max}/2}(\gamma))) \\
&\geqslant \int_{r_{\max}/2}^{r_{\max}} \mathrm{Length}(S_t \cap \partial\mathscr{N}_s(\gamma))\, ds \ ,
\end{aligned}$$

here $\mathrm{Area}(\cdot, g_t)$ means g_t-area and $\mathrm{Area}(\cdot)$ means hyperbolic area. There are two cases:

(i) if $S_t \cap \mathscr{N}_s(\gamma)$ has a component that is an annulus, then

$$\begin{aligned} \mathrm{Length}(S_t \cap \partial \mathscr{N}_s(\gamma)) &\geqslant \mathrm{Length}(\mathcal{A}_{\theta/l} \cap \partial \mathscr{N}_s(\gamma)) \\ &= 2(l^2 \cosh^2 s + \theta^2 \sinh^2 s)^{1/2} \\ &\geqslant 2\theta \sinh s \ , \end{aligned}$$

(ii) if $S_t \cap \mathscr{N}_s(\gamma)$ has a component that is a disk, then by assumption,

$$\mathrm{Length}(S_t \cap \partial \mathscr{N}_s(\gamma)) \geqslant 2\pi \sinh s \ .$$

When $\theta/l \gg 1$, so $r_{\max} \gg 1$ and $l \sinh r_{\max} \ll 1$, then we have the following estimates

$$\mathrm{Area}(S_t \cap \mathbb{T}(\gamma)) \geqslant 2c(\theta) \int_{r_{\max}/2}^{r_{\max}} \sinh s \, ds > c(\theta) \cosh r_{\max}$$

where $c(\theta) = \min\{\theta, \pi\}$, and

$$\mathrm{Area}(\partial \mathbb{T}(\gamma)) = 2\pi l \sinh r_{\max} \cosh r_{\max} < c(\theta) \cosh r_{\max} \ .$$

Recall that g_t metric is equal to the hyperbolic metric outside $\mathscr{N}_{\sigma(1-t)}(\gamma)$ and $\sigma(1-t) < \varepsilon$, thus we have

$$\mathrm{Area}(S_t \cap \mathbb{T}(\gamma), g_t) > \mathrm{Area}(\partial \mathbb{T}(\gamma), g_t) \ .$$

Now we can use cutting and pasting, i.e. replace each component of $S_t \cap \mathbb{T}(\gamma)$ by either an annulus or a disk on $\partial \mathbb{T}(\gamma)$, to get a new surface $S_t' \subset M \setminus \mathscr{N}_\varepsilon(\gamma)$ having the following properties

- $\mathrm{Area}(S_t', g_t) < \mathrm{Area}(S_t, g_t)$,
- $S_t' \cap \mathscr{N}_\varepsilon(\gamma) = \emptyset$, and
- S_t' is isotopic to S in $M \setminus \gamma$.

So S_t must be disjoint from $\mathscr{N}_\varepsilon(\gamma)$ for all sufficiently large t. □

By the Claim above, we see that S_t is a least area minimal surface with respect to the *hyperbolic metric* and $S_t \cap \mathscr{N}_\varepsilon(\gamma) = \emptyset$, where ε is a constant independent of t for all sufficiently large t, therefore all of them must be the same surface when t is sufficiently large. As $t \to 1$, we get a least area minimal surface T satisfying the conditions list in this theorem. □

Corollary 6.2. *Let $\Gamma = \{\gamma_i\}_{i=1}^n \subset M$ be the collection of mutually disjoint simple closed geodesics, with complex length $\mathscr{L}_i = l_i + \sqrt{-1}\,\theta_i$, such that $\theta_i/l_i \gg 1$ for $1 \leqslant i \leqslant n$, and let $S \subset M \setminus \Gamma$ be a closed embedded incompressible surface in M, then there exists a least area minimal surface $T \subset M \setminus \Gamma$ such that T is isotopic to S in $M \setminus \Gamma$.*

In particular, M contains at least 2^n minimal surfaces, which are incompressible, and which are not isotopic to each other in $M \setminus \Gamma$.

Proof. By Theorem 2.2, the tubes $\mathbb{T}(\gamma_1), \mathbb{T}(\gamma_2), \ldots, \mathbb{T}(\gamma_n)$ are mutually disjoint, so replace γ by Γ in the proof of Theorem 1.1, we can prove the first part of the corollary in the same way.

For the second part, J.-P. Otal proved in [**Ota03**, Theorem B] that Γ is *unlinked* in the following sense: There exists a homeomorphism between M and $\Sigma \times \mathbb{R}$ such that each component of Γ is contained in one of the surfaces $\Sigma \times i$, $1 \leqslant i \leqslant n$. Therefore the (incompressible) surfaces in $M \setminus \Gamma$ can separate Γ in

$$1 + \binom{n}{1} + \binom{n}{2} + \cdots + \binom{n}{n-1} + 1 = 2^n$$

ways such that they are not isotopic to each other in $M \setminus \Gamma$. By the first part, we have at least 2^n minimal surfaces. $\square$

References

[And83] Michael T. Anderson, *Complete minimal hypersurfaces in hyperbolic n-manifolds*, Comment. Math. Helv. **58** (1983), no. 2, 264–290.

[CG06] Danny Calegari and David Gabai, *Shrinkwrapping and the taming of hyperbolic 3-manifolds*, J. Amer. Math. Soc. **19** (2006), no. 2, 385–446.

[Mey87] Robert Meyerhoff, *A lower bound for the volume of hyperbolic* 3*-manifolds*, Canad. J. Math. **39** (1987), no. 5, 1038–1056.

[Mor82] Hiroshi Mori, *On surfaces of right helicoid type in H^3*, Bol. Soc. Brasil. Mat. **13** (1982), no. 2, 57–62.

[MSY82] William Meeks, III, Leon Simon, and Shing Tung Yau, *Embedded minimal surfaces, exotic spheres, and manifolds with positive Ricci curvature*, Ann. of Math. (2) **116** (1982), no. 3, 621–659.

[MY82a] William H. Meeks, III and Shing Tung Yau, *The classical Plateau problem and the topology of three-dimensional manifolds. The embedding of the solution given by Douglas-Morrey and an analytic proof of Dehn's lemma*, Topology **21** (1982), no. 4, 409–442.

[MY82b] William W. Meeks, III and Shing Tung Yau, *The existence of embedded minimal surfaces and the problem of uniqueness*, Math. Z. **179** (1982), no. 2, 151–168.

[Ota03] Jean-Pierre Otal, *Les géodésiques fermées d'une variété hyperbolique en tant que nœuds*, Kleinian groups and hyperbolic 3-manifolds (Warwick, 2001), London Math. Soc. Lecture Note Ser., vol. 299, Cambridge Univ. Press, Cambridge, 2003, pp. 95–104.

[SY79] R. Schoen and Shing Tung Yau, *Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature*, Ann. of Math. (2) **110** (1979), no. 1, 127–142.

[SY94] R. Schoen and S.-T. Yau, *Lectures on differential geometry*, International Press, Cambridge, MA, 1994.

[Tuz93] A. A. Tuzhilin, *Global properties of minimal surfaces in* $\mathbb{R}^3$ *and* $\mathbb{H}^3$ *and their Morse type indices*, Minimal surfaces, Adv. Soviet Math., vol. 15, Amer. Math. Soc., Providence, RI, 1993, pp. 193–233.

[Uhl83] Karen K. Uhlenbeck, *Closed minimal surfaces in hyperbolic* 3*-manifolds*, Seminar on minimal submanifolds, Ann. of Math. Stud., vol. 103, Princeton Univ. Press, Princeton, NJ, 1983, pp. 147–168.

[Wan08] Biao Wang, *Foliations for quasi-Fuchsian* 3*-manifolds*, 2008, available at `arXiv.org:0809.4057`.

Department of Mathematics
Cornell University
Ithaca, NY 14853-4201
E-mail address: wang@math.cornell.edu